

# LINEARIZED ANALYSIS OF THE DVALI-GABADADZE-PORRATI BRANE MODEL

Ioannis Giannakis and Hai-cang Ren <sup>†</sup>

*Physics Department, The Rockefeller University  
1230 York Avenue, New York, NY 10021-6399*

## Abstract

We solve the Einstein equations in the Dvali-Gabadadze-Porrati model with a static, spherically symmetric matter distribution on the *physical brane* and obtain an exact expression for the gravitational field outside the source to the first order in the gravitational coupling. Although, this expression when confined on the *physical brane* reproduces the correct  $4 - D$  Newtonian potential for distances  $r_s \ll r \ll \lambda$ , where  $\lambda$  is a characteristic length scale of the model, it does not coincide with the standard linearized form of the  $4 - D$  Schwarzschild metric. The solution reproduces the  $5 - D$  Schwarzschild metric in the linearized approximation for  $r \gg \lambda$ .

---

<sup>†</sup> e-mail: giannak@summit.rockefeller.edu, ren@summit.rockefeller.edu,

Recently there have been several proposals which attempt to achieve localization of gravity. The Randall-Sundrum model [1] is based on the assumption that ordinary matter and its gauge interactions are confined within a four-dimensional hypersurface embedded in a five-dimensional space which is *asymptotically anti-de Sitter*. Since gravity is formulated by five-dimensional general relativity, the gravitational field produced by a mass source on the brane might deviate from the standard Schwarzschild metric. Calculation of the form of the metric produced by a static, spherical mass distribution on the brane, to first order [2] and to second order [3], [4], in the gravitational coupling revealed no tangible disagreement with the classical tests of general relativity. The Randall-Sundrum model was shown also to be consistent with the gravitomagnetic effect [5]. Various aspects of black hole physics in the framework of the brane world have been discussed in [6].

An alternative gravity localized model, the Dvali-Gabadadze-Porrati model [7] consists of a four dimensional hypersurface embedded in a five-dimensional space which is *asymptotically Minkowski* ( for a recent review of brane models see [8]. ) The main motivation for this model is the competition between the bulk curvature scalar  $\mathcal{R}$  and the corresponding intrinsic curvature scalar  $R$  on the brane. The action which describes this particular model is given by

$$S = M_5^3 \int d^4x dy \sqrt{-g^{(5)}} \mathcal{R} + M_4^2 \int d^4x \sqrt{-g^{(4)}} R \quad (1)$$

where  $M_5$  stands for the 5 -  $D$  Planck constant and  $M_4$  is the 4 -  $D$  Planck constant.  $M_5$  and  $M_4$  are independent in general. The second term of (1) is included since matter localized on the brane might induce it through quantum fluctuations. This action yields the equations of motion

$$M_5^3 (\mathcal{R}_{mn} - \frac{1}{2} \mathcal{R} g_{mn}) + M_4^2 (R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu}) \delta_m^\mu \delta_n^\nu \delta(y) = \delta_m^\mu \delta_n^\nu T_{\mu\nu} \delta(y). \quad (2)$$

In this paper we shall solve the equations (2), with  $T_{\mu\nu} = M \eta_{\mu\nu} \delta^3(\vec{r})$  in the linearized approximation and we shall obtain an exact expression for the gravitational field outside the point source ( linearized analysis for the Randall-Sundrum model has been performed in [9].) For  $\vec{r} \neq 0$ , equation (2) implies

$$\mathcal{R} = -\frac{2}{3} \lambda R \delta(y), \quad (3)$$

and

$$\mathcal{R}_{mn} + \frac{\lambda}{3} g_{mn} R \delta(y) + \lambda (R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu}) \delta_m^\mu \delta_n^\nu \delta(y) = 0 \quad (4)$$

with  $\lambda = \frac{M_4^2}{M_5^3}$ , being a characteristic length scale of the model. Here and throughout the paper, we adopt the convention that the Greek indices take values 0-3 and the Latin indices 0-4. The interesting feature of this scenario is that although the spectrum of linearized fluctuations does not include a massless graviton bound state which would correspond to a

4 -  $D$  graviton it reproduces the correct Newtonian interaction on the brane for a certain range of parameters [10].

In order to describe the real world, the Dvali-Gabadadze-Porrati scenario has to satisfy all the existing tests of General Relativity. The most general axially symmetric and static metric in  $D = 4 + 1$  can always be brought to the following form

$$ds^2 = -e^a dt^2 + e^b dr^2 + e^c r^2 d\Omega^2 + dy^2, \quad (5)$$

where  $d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$  is the solid angle on  $S^2$  and  $a$ ,  $b$  and  $c$  are functions of  $r$  and  $y$ . The non-zero components of the 4 -  $D$  Ricci tensor for the general metric (5) in the linearized approximation are

$$\begin{aligned} R_{tt} &= -\frac{1}{2}a'' - \frac{1}{r}a' \\ R_{rr} &= \frac{1}{2}a'' + c'' - \frac{1}{r}b' + \frac{2}{r}c' \\ R_{\theta\theta} &= r^2 \left[ \frac{1}{2}c'' + \frac{2}{r}c' + \frac{a' - b'}{2r} \right] \\ R_{\phi\phi} &= R_{\theta\theta} \sin^2 \theta \end{aligned} \quad (6)$$

while the components of the 5 -  $D$  Ricci tensor read

$$\begin{aligned} \mathcal{R}_{tt} &= R_{tt} - \frac{1}{2}\ddot{a} \\ \mathcal{R}_{rr} &= R_{rr} + \frac{1}{2}\ddot{b} \\ \mathcal{R}_{\theta\theta} &= R_{\theta\theta} + \frac{r^2}{2}\ddot{c} \\ \mathcal{R}_{\phi\phi} &= \mathcal{R}_{\theta\theta} \sin^2 \theta \\ \mathcal{R}_{yy} &= \frac{1}{2}(\ddot{a} + \ddot{b} + 2\ddot{c}) \\ \mathcal{R}_{ry} &= \mathcal{R}_{yr} = \frac{1}{2} \left[ \dot{a}' + 2\dot{c}' - \frac{2}{r}(\dot{b} - \dot{c}) \right] \end{aligned} \quad (7)$$

where  $a'$  and  $\dot{a}$  indicate differentiation with respect to  $r$  and  $y$  respectively. Consequently the 4 -  $D$  and the 5 -  $D$  Ricci scalars read

$$\begin{aligned} R &= a'' + 2c'' + \frac{2}{r}a' - \frac{2}{r}b' + \frac{6}{r}c' + \frac{2}{r^2}(b + c) \\ \mathcal{R} &= R + \ddot{a} + \ddot{b} + 2\ddot{c}. \end{aligned} \quad (8)$$

These equations apply to the positive side of the brane, i.e.,  $y > 0$ , the corresponding equations to the negative side of the brane,  $y < 0$ , are obtained by  $Z_2$  symmetry. The

Einstein equations (3) and (4) then demand that all of the components of  $\mathcal{R}_{mn}$  vanish for  $y \neq 0$  and dictate the following matching conditions at  $y = 0$

$$\begin{aligned} \dot{a}(r, y)|_{y=0} &= -\frac{\lambda}{2}(a'' + \frac{2}{r}a')|_{y=0} + \frac{\lambda}{6}R|_{y=0} \\ \dot{b}(r, y)|_{y=0} &= -\frac{\lambda}{2}(a'' + 2c'' - \frac{2}{r}b' + \frac{4}{r}c')|_{y=0} + \frac{\lambda}{6}R|_{y=0} \\ \dot{c}(r, y)|_{y=0} &= -\frac{\lambda r^2}{2}(c'' + \frac{4}{r}b' + \frac{a' - b'}{r})|_{y=0} + \frac{\lambda}{6}R|_{y=0}. \end{aligned} \quad (9)$$

The  $yy$  component of the equation (4), together with equation (3) implies that  $R = 0$  for  $y \neq 0$ , which leads to  $R = 0$  at  $y = 0$  by continuity. This simplifies the matching conditions (9)

$$\begin{aligned} \dot{a}(r, y)|_{y=0} &= -\frac{\lambda}{2}(a'' + \frac{2}{r}a')|_{y=0} = \frac{\lambda}{2}\ddot{a}|_{y=0} \\ \dot{b}(r, y)|_{y=0} &= -\frac{\lambda}{2}(a'' + 2c'' - \frac{2}{r}b' + \frac{4}{r}c')|_{y=0} = \frac{\lambda}{2}\ddot{b}|_{y=0} \\ \dot{c}(r, y)|_{y=0} &= -\frac{\lambda r^2}{2}(c'' + \frac{4}{r}b' + \frac{a' - b'}{r})|_{y=0} = \frac{\lambda}{2}\ddot{c}|_{y=0}. \end{aligned} \quad (10)$$

The solution to the equation  $\mathcal{R}_{tt} = 0$  reads

$$a(r, y) = \frac{1}{2\pi^2} \int_0^\infty dp p^2 C(p) j_0(pr) e^{-py}, \quad y > 0, \quad (11)$$

where  $C(p)$  is a function of  $p$  to be determined and  $j_0(x)$  is the spherical Bessel function. Substituting (11) into the first of equations (9) and taking into account that  $\delta^3(\vec{r}) = \int \frac{d^3\vec{p}}{(2\pi)^3} e^{i\vec{p}\vec{r}} = \frac{1}{2\pi^2} \int_0^\infty dp p^2 j_0(pr)$  we find

$$C(p) = \frac{C_0}{p + \frac{1}{2}\lambda p^2}. \quad (12)$$

As a result  $a(r, y)$  is given by the expression

$$a(r, y) = \frac{C_0}{2\pi^2} \int_0^\infty dp \frac{p^2}{p + \frac{1}{2}\lambda p^2} j_0(pr) e^{-py}, \quad y > 0, \quad (13)$$

where  $C_0$  is a constant to be determined by the requirement that the solution reproduces the correct Newtonian limit. The Newtonian limit is specified by the asymptotic behavior of  $g_{00}$  on the brane for large  $r$ . Since  $p \sim \frac{1}{r}$  and  $p + \frac{\lambda}{2}p^2 \sim \frac{1}{r} + \frac{\lambda}{2r^2}$  we find that for  $r \ll \lambda$

$$a(r, y=0) \simeq \frac{C_0}{\pi^2 \lambda} \int_0^\infty dp j_0(pr) = \frac{C_0}{2\pi \lambda r} = -\frac{2GM}{r} \quad (14)$$

which determines the relation  $C_0 = -4\pi GM\lambda$ .

The  $yy$ -component of the equations of motion outside the brane provide the equation  $\ddot{a} + \ddot{b} + 2\ddot{c} = 0$ . If we impose the following conditions to the solution

$$\lim_{|y|, r \rightarrow \infty} a = \lim_{|y|, r \rightarrow \infty} b = \lim_{|y|, r \rightarrow \infty} c = 0 \quad (15)$$

independent of the order of the limits, we find that

$$a + b + 2c = 0, \quad (16)$$

which implies  $R = 0$  on the brane. By combining equations  $\mathcal{R}_{ry} = 0$  and (16) we derive the following expression for  $b(r, y)$

$$b(r, y) = -\frac{1}{r^3} \int^r dr' r'^2 a(r', y). \quad (17)$$

Consequently by substituting (13) into (17) we find

$$b(r, y) = \frac{2GM\lambda}{\pi} \int_0^\infty dp \frac{p^2}{p + \frac{1}{2}\lambda p^2} \frac{j_1(pr)}{pr} e^{-py}, \quad y > 0. \quad (18)$$

Finally equation (16) provides an expression for  $c(r, y)$

$$c(r, y) = \frac{GM\lambda}{\pi} \int_0^\infty dp \frac{p^2}{p + \frac{1}{2}\lambda p^2} [j_0(pr) - \frac{j_1(pr)}{pr}] e^{-py}, \quad y > 0. \quad (19)$$

Although  $a(r, y)$  satisfies the matching condition,  $b(r, y)$  and  $c(r, y)$  fail to do so. The boundary condition for  $b(r, y)$  is (10)

$$\dot{b}(r, y)|_{y=0} = -\lambda \left( \frac{1}{2} a'' + c'' - \frac{1}{r} b' + \frac{2}{r} c' \right) |_{y=0} = \frac{\lambda}{2} \ddot{b}(r, y)|_{y=0}. \quad (20)$$

By substituting equation (18) into (20) we find

$$\frac{\lambda}{2} \ddot{b}(r, y)|_{y=0} - \dot{b}(r, y)|_{y=0} = \frac{2GM\lambda}{\pi} \int_0^\infty dp p^2 \frac{j_1(pr)}{pr} = \frac{GM\lambda}{r^3} \quad (21)$$

which is nonzero. The situation is reminiscent with the Randall-Sundrum model where two sets of solutions to the Einstein equations in two different coordinate systems, or equivalently, in two different gauges were constructed. The solution that we found is expressed in coordinates which are straight with respect to the horizon and is free from metric singularities far away from the source but it causes the brane to bend away from  $y = 0$  [11]. By transforming the solution to coordinates based on the brane we shall obtain another solution which satisfies the matching condition. In this set of coordinates the brane appears straight. We shall denote the solution in this gauge by  $a^P(r, y)$ ,  $b^P(r, y)$  and  $c^P(r, y)$ , where the superscript  $P$  means *physical*.

We shall now discuss the gauge symmetries of the Einstein equations, more specifically the coordinate transformations that respect the axially symmetric, static form of the five dimensional metric (5). Let's perform a coordinate transformation (gauge transformation) generated by  $v, u$ , functions of  $r$  and  $y$  such that  $r \mapsto r + v(r, y)$  and  $y \mapsto y + u(r, y)$  and demand that this particular coordinate transformation respects the form of the five dimensional metric (5). We find that the functions  $v, u$  obey to linear order the following relations

$$\dot{v} = 0, \quad v' + \dot{u} = 0, \quad (22)$$

while the components of the metric transform under these residual coordinate transformations as follows

$$\delta a = 0, \quad \delta b = 2u', \quad \delta c = 2\frac{u}{r}. \quad (23)$$

We can easily verify that the Einstein equations remain invariant under the transformations (22). Thus, the physical solution will be given by

$$a^P = a + \delta a, \quad b^P = b + \delta b, \quad c^P = c + \delta c. \quad (24)$$

We impose the boundary condition on  $b^P$  on the brane,

$$\frac{\lambda}{2} \ddot{b}^P(r, y)|_{y=0} - \dot{b}^P(r, y)|_{y=0} = 0. \quad (25)$$

Under a coordinate transformation generated by  $u, v$  we find that

$$\lambda \ddot{u}'(r, 0) - 2\dot{u}'(r, 0) = -\frac{GM\lambda}{r^3}. \quad (26)$$

The parameters of the transformation,  $u$  and  $v$ , satisfy equations (22). Consequently,

$$v = v(r), \quad u(r, y) = -v'(r)y + \chi(r), \quad (27)$$

where  $\chi(r)$  is an arbitrary function of  $r$ . By substituting (27) into (26) we find that

$$v(r) = -\frac{GM\lambda}{4r}, \quad u(r, y) = -\frac{GM\lambda}{4r^2}y + \chi(r). \quad (28)$$

The physical solution to the first order in the gravitational coupling takes then the form

$$\begin{aligned} a^P(r, y) &= -\frac{2GM\lambda}{\pi} \int_0^\infty dp \frac{p^2}{p + \frac{1}{2}\lambda p^2} j_0(pr) e^{-py} \\ b^P(r, y) &= \frac{2GM\lambda}{\pi} \int_0^\infty dp \frac{p^2}{p + \frac{1}{2}\lambda p^2} \frac{j_1(pr)}{pr} e^{-py} + \frac{GM\lambda}{r^3} y + 2\chi'(r) \\ c^P(r, y) &= \frac{GM\lambda}{\pi} \int_0^\infty dp \frac{p^2}{p + \frac{1}{2}\lambda p^2} [j_0(pr) - \frac{j_1(pr)}{pr}] e^{-py} - \frac{GM\lambda}{2r^3} y + \frac{2\chi(r)}{r}. \end{aligned} \quad (29)$$

Finally for  $2GM \ll r, y \ll \lambda$  the solution takes the form

$$\begin{aligned} a(r, y) &= -\frac{2GM}{i\pi r} \ln \frac{y + ir}{y - ir} = -\frac{4GM}{\pi r} \tan^{-1} \frac{r}{y} = -\frac{2GM}{r} \left(1 - \frac{2}{\pi} \tan^{-1} \frac{r}{y}\right) \\ b(r, y) &\cong \frac{GM}{r} \left[ \left(1 - \frac{y^2}{r^2}\right) \left(1 - \frac{2}{\pi} \tan^{-1} \frac{r}{y}\right) + \frac{2y}{\pi r} \right] \\ c(r, y) &= -\frac{1}{2}(a + b) \cong \frac{GM}{2r} \left[ \left(1 + \frac{y^2}{r^2}\right) \left(1 - \frac{2}{\pi} \tan^{-1} \frac{r}{y}\right) - \frac{2y}{\pi r} \right] \end{aligned} \quad (30)$$

We observe that if we confine the metric on the brane  $y = 0$ , the *physical solution* reads for  $2GM \ll r \ll \lambda$

$$a^P(r) = -\frac{2GM}{r}, \quad b^P(r) = \frac{GM}{r} + 2\chi', \quad c^P(r) = \frac{GM}{2r} + \frac{2\chi}{r}, \quad (31)$$

where  $\chi$  is an arbitrary function of  $r$ . In order to recover the standard static, spherically symmetric metric on the brane, i.e.  $c^P = 0$  at  $y = 0$  we choose  $\chi = -\frac{GM}{4}$ . This leaves  $b^P = \frac{GM}{r}$ , which differs from the standard form of the Schwarzschild metric and thus leads to different predictions for the bending of light. In the Randall-Sundrum model the choice of  $\chi$  that sets  $c^P = 0$  leads also to  $b^P = \frac{GM}{2r}$  which is the form of the Schwarzschild metric in the linearized approximation.

Similarly for  $r, y \gg \lambda$

$$\begin{aligned} a(r, y) &\cong -\frac{2GM\lambda}{r^2 + y^2} \\ b(r, y) &\cong \frac{2GM\lambda}{r^2} \left(1 - \frac{y}{r} \tan^{-1} \frac{r}{y}\right) \\ c(r, y) &= -\frac{1}{2}(a + b) \cong \frac{GM\lambda}{\pi r^2} \left(\frac{y^2}{y^2 + r^2} - \frac{y}{r} \tan^{-1} \frac{r}{y}\right). \end{aligned} \quad (32)$$

We note that if we confine the metric on the *physical brane*  $y = 0$ , the solution reads

$$a^P(r) = -\frac{2GM\lambda}{r^2}, \quad b^P(r) = \frac{2GM\lambda}{r^2}, \quad c^P(r) = 0, \quad (33)$$

which agrees with the standard linearized form of the 5 -  $D$  Schwarzschild metric.

In this paragraph we shall recapitulate what we have done in this paper. We solved the linearized Einstein equations in the model proposed by Dvali, Gabadadze and Porrati with a static, spherically symmetric matter distribution on the *physical brane* and found an exact expression for the gravitational field. Although when confined on the *physical brane* the metric reproduces the 4 -  $D$  Newtonian potential for a certain range of parameters, more specifically for  $2GM \ll r \ll \lambda$  it fails to reproduce the standard Schwarzschild metric in its linearized form. Does this result then imply that the model is incompatible

with the experimental predictions of general relativity? Recently in [12] it was argued that the linearized approximation ceases to be reliable for this range of parameters and subsequently an exact classical solution to all orders in the gravitational coupling is needed. This argument is based on an early work by Vainshtein [13] in which he suggested that the limit of vanishing mass of massive general relativity may coincide with the predictions of massless general relativity in contrast with previous papers [14], [15] if the nonperturbative exact classical solution is used. The analog of the graviton mass on the brane is the inverse of the length scale  $\lambda$ . The possibility then remains that a nonperturbative solution exists and it reproduces the standard Schwarzschild solution when confined on the *physical brane*. On the contrary the metric for  $r \gg \lambda$  behaves as a  $5 - D$  Schwarzschild metric. (See also [16] where it is claimed that four-dimensional behavior is recovered at large distances in the Dvali-Gabadadze-Porrati model.) We hope to report in the near future results which deal with these issues.

Beyond the linear approximation, the properties of an event horizon which we established in a rigorous manner for the Randall-Sundrum model [17] apply to the DGP brane model as well. If a Schwarzschild black hole localized on the brane exists, the extension of the horizon into the bulk with the Gauss normal form of the metric (5) is tubular. This is not in contradiction with the analysis based on the linearized approximation, employed in this work, since the approximation breaks down for  $\lambda y \gg r^2$ , thus permitting the horizon to extend to  $y \mapsto \infty$ .

## Acknowledgments.

We would like also to thank A. Polychronakos, M. Porrati and G. Siopsis for useful discussions. This work was supported in part by the Department of Energy under Contract Number DE-FG02-91ER40651-TASK B.

## References.

- [1] L. Randall and R. Sundrum, Phys. Rev. Lett. **83** (1999) 4690.
- [2] S. Giddings, E. Katz and L. Randall, JHEP **0003** (2000) 023.
- [3] I. Giannakis and H. C. Ren, Phys. Rev. **D63** (2001) 024001.
- [4] H. Kudoh and T. Tanaka, Phys. Rev. **D64** (2001) 084022.
- [5] A. Nayeri and A. Reynolds, “ *Gravitomagnetism in Brane Worlds* ”, hep-th/0107201.
- [6] A. Chamblin, S. W. Hawking and H. S. Reall, Phys. Rev. **D61** (2000) 065007, T. Shiromizu and M. Shibata, Phys. Rev. **D62** (2000) 127502, N. Dadhich, R. Maartens, P. Papadopoulos and V. Rezania, Phys. Lett. **487B** (2000) 1, A. Chamblin, H. Reall, H. Shinkai and T. Shiromizu, Phys. Rev. **D63** (2001) 064015, R. Emparan, G. Horowitz and R. Myers, JHEP **0001** (2000) 007; Phys. Rev. Lett. **85** (2000) 499, J.



- Garriga and M. Sasaki, Phys. Rev. **D61** (2000) 065007, I. Giannakis and H. c. Ren, Phys. Rev. **D64** (2001) 065015, I. Oda, “*Reissner-Nordstrom Black Hole in Gravity Localized Models*”, hep-th/0008055, M. Bruni, C. Germani and R. Maartens, “*Gravitational Collapse on the Brane*”, gr-qc/0108013, S. Vacaru and E. Gabunov, “*Anisotropic Black Holes in Einstein and Brane Gravity*”, hep-th/0108065, M. S. Modgil, S. Panda and G. Sengupta, “*Rotating Brane World Black Holes*”, hep-th/0104122, W. T. Kim, J. Oh, M. K. Oh and M. S. Yoon, “*Brane World Black Holes in Randall-Sundrum models*”, hep-th/0006134, P. Kanti and K. Tamvakis, “*Quest for Localized 4-D Black Holes in Brane Worlds*”, hep-th/0110298, T. Wiseman, “*Relativistic Stars in Randall-Sundrum Gravity*”, hep-th/0111057.
- [7] G. Dvali, G. Gabadadze and M. Porrati Phys. Lett. **485B** (2000) 208.
  - [8] R. Dick, Class. Quant. Grav. **18** (2001) R1.
  - [9] I. Y. Arefe’va, M. Ivanov, W. Muck, K. S. Viswanathan, and I. V. Volovich, Nucl. Phys. **B590** (2000) 273; Y. S. Myung, G. Kang and H. W. Lee, Phys. Lett. **478B** (2000) 294; H. Collins and B. Holdom, Phys. Rev. **D62** (2000) 124008; W. Muck, K. S. Viswanathan and I. V. Volovich, Phys. Rev. **D62** (2000) 105019; R. Dick and D. Mikulovicz, Phys. Lett. **476B** (2000) 363.
  - [10] G. Dvali, G. Gabadadze and M. Porrati, Phys. Lett. **484B** (2000) 112.
  - [11] J. Garriga and T. Tanaka, Phys. Rev. Lett. **84** (2000) 2778.
  - [12] C. Deffayet, G. Dvali, G. Gabadadze and A. Vainshtein, “*Nonperturbative Continuity in Graviton Mass versus Perturbative Discontinuity*”, hep-th/0106001.
  - [13] A. Vainshtein, Phys. Lett. **39B** (1972) 393.
  - [14] H. van Dam and M. Veltman, Nucl. Phys. **B22** (1970) 397
  - [15] V. I. Zakharov, JETP Lett. **12** (1970) 312.
  - [16] J. Avery, R. Mahurin and G. Siopsis, “*A Brane in five-dimensional infinite flat Minkowski space*”, hep-th/0108132.
  - [17] I. Giannakis and H. c. Ren, Phys. Rev. **D63** (2001) 125017.